

A Perturbation Method for the 3D Finite Element Modeling of Electrostatically Driven MEMS



M. Boutaayamou, R. V. Sabariego, and P. Dular




Department of Electrical Engineering and Computer Science (ELAP) University of Liège, Belgium

Modeling electrostatically driven MEMS

- Lumped spring-mass models
 - helpful for physical insight
 - neglect important effects such as bending of plates and the fringing field effects
- Finite element (FE) method
 - adapted for complex geometries
 - compute accurately the fringing field effects at the expense of dense discretization near corners
 - movement modeling usually requires successive meshing and computations for each new position
 - classical approach computationally **expensive**
- Perturbation method
 - **unperturbed field** computed in a global domain without the presence of some conductive regions
 - ✓ **no mesh** of perturbing domains
 - ✓ possible symmetries or analytical solution
 - ✓ applied source to the perturbation problems
 - **perturbed field** determined in a local domain
 - ✓ **subproblem-adapted meshes**
- Iterative sequence of perturbation problems
 - when coupling between sub-regions is significant
 - ✓ **successive perturbations** in each region are computed from one region to the other
 - ✓ each sub-problem gives a **correction** as a perturbation

Perturbation method


- **Unperturbed domain:** parallel-plate capacitor
- **Perturbing domain:** micro-beam (100μm × 10μm)



Unperturbed problem

- Unperturbed electric scalar potential FE formulation

$$(-\epsilon \text{grad } v_u, \text{grad } v')_{\Omega} - \langle \mathbf{n} \cdot \mathbf{d}_u, v' \rangle_{\Gamma_u} = 0$$



$\text{grad } v_u$ is projected on $\partial\Omega_{c,p}$...

Perturbation equations


- Perturbed electric scalar potential FE formulation

$$\langle \text{grad } v_s, \text{grad } v' \rangle_{\partial\Omega_{c,p}} - \langle \text{grad } v_u, \text{grad } v' \rangle_{\partial\Omega_{c,p}} = 0$$


$$(-\epsilon \text{grad } v, \text{grad } v')_{\Omega_p} - \langle \mathbf{n} \cdot \mathbf{d}_p, v' \rangle_{\Gamma_p} = 0$$

$$v = -v_s|_{\partial\Omega_{c,p}}$$


...with $v_u + v = v_p$ & $e_u + e = e_p$






Post-processing




Iterative process of perturbation problems





Application



$d_1 = 3 \mu\text{m}$



$d_1 = 3 \mu\text{m}$



Tolerance = 1% (with respect to the regular FEM)

Conclusions

- A **perturbation** approach based on electric scalar potential FE formulation has been presented
 - adapted for complex geometries
 - unperturbed field computed in a global domain **without the presence of conductive regions**
 - unperturbed field applied as **source** in $\Omega_{c,p}$
 - perturbed field determined in **reduced domain** Ω_p
 - **iterative procedure** used when the coupling between the sub-regions is significant
- Convergence acceleration of iterative sequence in progress